

Layer (heights managed by coppicing)	Species used in Trial Woody Meadow	substitutes
<b>Base (&lt;1 m height) - 70-75% of planting</b>		
	<i>Commersonia hermannifolia</i>	<i>Kennedia prostrata</i> / <i>Myoporum parvifolium</i> / <i>Commersonia hermannifolia</i>
	<i>Scaevola aemula</i>	<i>Scaevola aemula</i> / <i>Scaevola albida</i> / <i>Mentha australis</i>
	<i>Correa reflexa</i>	<i>Correa alba</i> / <i>Correa decumbens</i> / <i>Melaleuca thymifolia</i>
	<i>Goodenia ovata</i> Prostrate	<i>Brachyscome multifida</i> / <i>Coronidium scorpioides</i>
	<i>Lasiopetalum ferrugineum</i>	<i>Veronica perfoliata</i> / <i>Bracteantha bracteata</i> 'Diamond Head'
	<i>Veronica perfoliata</i>	<i>Zieria prostrata</i> / <i>Austromyrtus dulcis</i> / <i>Gonocarpus teucroides</i> / <i>Commersonia hermannifolia</i> <i>Opercularia</i> / <i>Pomax</i>
<b>Middle (1-2 m height) - 20-25% of planting</b>		
	<i>Leptospermum polygalifolium</i> subsp. <i>polygalifolium</i>	<i>Leptospermum arachnoides</i> / <i>Gompholobium latifolium</i> / <i>Acacia myrtifolia</i> / <i>Zieria smithii</i>
	<i>Platylobium formosum</i>	<i>Pultenaea retusa</i> / <i>Cassinia</i> species
	<i>Indigofera australis</i>	<i>Acacia echinula</i> / <i>Goodenia ovata</i>
	<i>Prostanthera denticulata</i>	<i>Westringia wynyabbie gem</i> / <i>Prostanthera denticulata</i>
	<i>Banksia spinulosa</i>	<i>Bauera rubioides</i> / <i>Leptospermum rotundifolium</i>
	<i>Pultenaea retusa</i>	
<b>Upper (&gt;2 m height) - 0-5% of planting</b>		
	<i>Acacia stricta</i>	<i>Callistemon pinifolius</i> / <i>Banksia oblongifolia</i>
	<i>Callistemon citrinus</i>	<i>Kunzea ambigua</i>
	<i>Gompholobium latifolium</i>	
	<i>Ozothamnus diosmifolius</i>	<i>Olearia phlogopappa</i> / <i>Babingtonia pluriflora</i>